


Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

## International Journal of Gerontology

journal homepage: [www.ijge-online.com](http://www.ijge-online.com)

## Original Article

Changes in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Regulatory T Cells in Relation to Aging and Lung Tumor Incidence<sup>☆</sup>Li-Jing Zhu<sup>1</sup>, Pan-Fei Hou<sup>2</sup>, Ling Wang<sup>1\*</sup>, Guang-Bo Zhang<sup>1</sup>, Yan Xie<sup>1</sup>, Xu-Dong Pan<sup>1</sup>, Ting-Ting Chang<sup>1</sup><sup>1</sup>Clinical Immunology Laboratory, The First Affiliated Hospital of Soochow University, Jiangsu, <sup>2</sup>Rushan Hospital of Binzhou Medical University, Shandong, China

## ARTICLE INFO

## Article history:

Received 1 November 2010

Received in revised form

23 February 2011

Accepted 17 March 2011

Available online 20 March 2012

## Keywords:

aging,  
lung tumor,  
regulatory T cells,  
relation

## SUMMARY

**Background:** CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> regulatory T (Treg) cells mediate immunosuppression and play an important role in tumor immune evasion. Studies have demonstrated that this cell population represents an aging-related change. It is not clear whether this change leads to higher tumor incidence in the elderly. We investigated changes in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells in relation to aging and tumor incidence.**Methods:** We set up a Lewis lung cancer model with 26 C57BL/6 female mice. The animals were divided into six groups: young healthy, middle-aged healthy, elderly healthy, young tumor, middle-aged tumor and elderly tumor. We evaluated changes in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup>Treg cells in the spleen of all animals using a flow cytometry method. Levels of Foxp3 mRNA in splenocytes were measured using a real-time RT-PCR method. **Results:** The CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup>/CD4<sup>+</sup> T cell ratio ( $t = 2.23$ ,  $p = 0.032$ ) and Foxp3 mRNA levels ( $t = 3.26$ ,  $p = 0.0042$ ) were higher in the tumor groups than in the healthy groups. In the healthy groups, there was a significant increase in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells on aging ( $F = 47.70$ ,  $p = 0.000$ ); elderly mice had a significantly greater population of CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells in spleen compared to the younger groups. The highest population was observed in the elderly tumor group. The same trend was evident for Foxp3 mRNA ( $F = 6.56$ ,  $p = 0.0090$ ).**Conclusions:** The results suggest a close relationship between changes in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells and aging and lung tumor genesis and development.

Copyright © 2012, Taiwan Society of Geriatric Emergency &amp; Critical Care Medicine. Published by Elsevier Taiwan LLC. All rights reserved.

## 1. Introduction

With aging, the incidence and prevalence of tumor increase<sup>1–3</sup>, suggesting an intimate relationship between aging and tumors<sup>4,5</sup>. In other words, many mechanisms involved in the aging process share molecular mechanisms implicated in carcinogenesis<sup>6</sup>. The existence of relationships between aging and the risk of tumor development is not very well understood. From an immunology perspective, a tumor suppresses the immune response of the body and develops *in vivo* via many pathological processes, such as immunity escape, toleration and suppression<sup>7</sup>. Changes in the immunity system with aging further provide favorable conditions for tumor escape, so there is a higher incidence of tumors in the elderly. An understanding of the contribution of immunosenescence to tumor development and progression may help in

designing better interventions for tumors in the elderly. Recent evidence demonstrated that regulatory T cell (Treg)-mediated immunosuppression is a crucial tumor immune evasion mechanism and contributes to the failure of tumor immunotherapy<sup>8–11</sup>. This regulatory capacities has been attributed to several different cell types, but the most characterized group is CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells, which increase with aging<sup>12</sup>. We presume that increases in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells are responsible for the high tumor incidence in the elderly. Recently, lung tumor, as the highest rate of the malignant tumors<sup>13</sup>, rises exponentially with aging<sup>14</sup>. In this study, we set up a Lewis lung cancer model in animals of different ages. We investigated changes in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells in healthy and tumor model groups with aging and explored the relation between immunosenescence and lung tumors.

## 2. Materials and methods

## 2.1. Cell lines

Transplantable Lewis lung carcinoma (LLC) cells (Cell Bank, Shanghai Scientific Research Institute, Chinese Academy of Sciences)

<sup>☆</sup> All contributing authors declare no conflicts of interest.

\* Correspondence to: Dr Ling Wang, The First Affiliated Hospital of Soochow University, Suzhou 215006, Jiangsu Province, China.

E-mail address: [wangling40@126.com](mailto:wangling40@126.com) (L. Wang).

were cultivated *in vitro* in DMEM culture medium with a high glucose content supplemented with 10% fetal bovine serum, 100 U/mL penicillin, 100 µg/mL streptomycin sulfate, and 0.25 µg/mL amphotericin B in a 5% CO<sub>2</sub> atmosphere at 37 °C.

## 2.2. Animals and tumor model

C57BL/6 female mice ( $n = 36$ , Institute of Laboratory Animals, Shanghai SLAK, Certificate of Quality No. 2007000504834) were divided into six groups: 4-week healthy (young healthy, YH), 6-month healthy (middle-aged healthy, MH), 20-month healthy (elderly healthy, EH), 4-week tumor (young tumor, YT), 6-month tumor (middle-aged tumor, MT) and 20-month tumor (elderly tumor, ET), with six mice in each group. Mice were maintained under specific pathogen-free conditions and provided with food and water *ad libitum*. All animals were housed for 1 week and maintained under standard conditions prior to experimentation. All animal procedures were performed in accordance with the rules of the local Ethics Committee and the guidelines established by the Animal Care Committee, which approved the study (No. 2091116). All tumor groups were injected subcutaneously with 0.1 mL of LLC cells ( $2 \times 10^6$  cells) in the left forelimb. The healthy groups were injected with the same amount of phosphate-buffered saline (PBS) in the same site. After implantation, tumor growth dynamics was monitored every 2 days. The tumor long diameter ( $L$ ) and short diameter ( $W$ ) were used to calculate tumor volume according to the Steel formula:  $V = LW^2/2$ . Tumor growth curves were plotted for all experimental groups.

## 2.3. Flow cytometry

The mice were sacrificed 20 days after the injection treatments. The spleen was harvested and a splenocyte suspension was prepared. Lymphocytes were isolated from this suspension by density centrifugation over a Ficoll-Hypaque gradient (Lymphoprep, Tian Jin Hao Yang Biological Manufacture, Tianjin, China). An aliquot of 100 µL ( $1 \times 10^6$  cells) was removed for dyeing, and 1 mL of RNAiso Plus was added to the remainder for preservation at  $-80^\circ\text{C}$  for measurement of *Foxp3* mRNA.

A Mouse Regulatory T cell staining kit (ebioscience, San Diego, USA) was used for staining according to the manufacturer's directions. In brief, 100 µL of prepared cells ( $1 \times 10^6$ ) was added to individual tubes and surface molecules were stained using a standard procedure, with a fluorescein isothiocyanate-CD4 antibody and an allophycocyanin-conjugated CD25 antibody. After washing with flow cytometry staining buffer, cells were treated with fixation/permeabilization working solution and washed once with 2 mL of permeabilization buffer. After centrifugation, the supernatant was decanted and cells were then stained with a phycoerythrin-conjugated *Foxp3* antibody.

Finally, cells were washed with 2 mL of permeabilization buffer, resuspended in flow cytometry staining buffer, and analyzed by flow cytometry (Beckman Coulter, Inc. California, USA).

## 2.4. Real-time RT-PCR

Spleen cells were isolated and preserved at  $-80^\circ\text{C}$ . DNA-free RNA templates were prepared using an RNA isolation kit (Shingene Bio-Technologies, Shanghai, China). RNA concentrations and quality were assessed using spectrophotometry at wavelengths of 260 and 280 nm. RNA was reverse-transcribed using a reverse transcription kit (Takara Biotechnology, Dalian, China) according to the manufacturer's instructions. Real-time PCR assays were carried out using a SYBR® Premix Ex Taq™ kit (Takara Biotechnology). The 20-µL reaction system contained SYBR Premix Ex Taq (10 µL), ROX Reference Dye II (0.4 µL), *Foxp3* primers (5'-CAG CTG CCT ACA GTG CCC CTA G-3' and 5'-CAT TTG CCA GCA GTG GGT AG-3', 0.4 µL each) cDNA (2 µL) and dH<sub>2</sub>O (6.8 µL). PCR was performed on an ABI 7500 real-time PCR system (Applied Biosystems, Inc. California, USA). The conditions were 95 °C for 30 s, followed by 40 cycles at 95 °C for 5 s and 60 °C for 34 s. HPRT (5'-GTT GGA TAC AGG CCA GAC TTT GTT G-3' and 5'-GAT TCA ACT TGC TCT CAT CTT AGG C-3') was used as a housekeeping gene for normalization of gene transcript levels.

## 2.5. Statistical analysis

Values are presented as mean  $\pm$  standard deviation (SD). Statistical analysis was performed using SAS version 8.0 software. Student's *t* test was used to examine the significance of differences between two groups. One-way analysis of variance (ANOVA), followed by Dunnett's test and a *q* test, was used to compare means across multiple groups. A *p* value of  $< 0.05$  was deemed to be statistically significant.

## 3. Results

### 3.1. Establishment of the tumor model

We applied a classic approach to establish a lung tumor model. After injection of LLC cells, tumors were palpable on the 10<sup>th</sup> day and could be seen by the naked eye on the 16<sup>th</sup> day. Tumor growth curves are shown in Fig. 1. By contrast, the healthy groups showed no changes. All the mice were killed 20 days after implantation (Fig. 2) and splenocytes were isolated and used in experiments.

### 3.2. Changes in CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cells in spleen

The presence of CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cells was evaluated in spleen for the six groups. As shown in Fig. 3 and Table 1, the ratio




Fig. 1. Tumor growth curves for the experimental groups.



**Fig. 2.** A 4-week-old mouse sacrificed 20 days after injection. Tumor tissue in the left forelimb is clearly evident.

of CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> to CD4<sup>+</sup> T cells was higher in the tumor groups than in the healthy groups ( $t = 2.23$ ,  $p = 0.032$ ). The ratio showed significant differences among the healthy groups ( $F = 47.70$ ,  $p = 0.000$ ). The MH group had a larger population of CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells in spleen compared to the YH group ( $F = 53.35$ ,  $p = 0.0019$ ) and there was a significant difference between the EH and MH groups ( $F = 22.02$ ,  $p = 0.0082$ ). The highest ratio was observed for the ET group.

**Table 1**  
Comparison of the CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup>/CD4<sup>+</sup> cell ratio in spleen for the six groups.

Group	Healthy	Tumor
Young (4 w)	1.56 ± 0.43	2.13 ± 0.33
Middle-aged (6 m)	2.60 ± 0.30*	3.25 ± 0.42
Elderly (20 m)	3.71 ± 0.40**	5.01 ± 0.24
<i>F</i> values	47.70	108.29
<i>P</i> values	<0.01	<0.01


\*  $p = 0.0019$  compared with the young healthy, \*\*  $p = 0.0082$  compared with the middle-aged healthy group.

3.3. *Foxp3* mRNA expression levels in spleen

We investigated the levels of *Foxp3* mRNA for the six groups. *Foxp3* mRNA amplification was successfully performed by RT-PCR (Fig. 4). Our results revealed differences among the six groups (Fig. 5). *Foxp3* mRNA levels were higher in the tumor compared to the healthy groups ( $t = 3.26$ ,  $p = 0.0042$ ). *Foxp3* mRNA levels in spleen also increased with aging ( $F = 6.56$ ,  $p = 0.0090$ ); among the healthy groups, levels were significantly lower in the YH group compared to the MH and EH groups. In agreement with its immune function, changes in *Foxp3* expression were consistent with variations in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells. Levels of *Foxp3* mRNA were affected by both age and tumor incidence, and the highest levels were found in the elderly tumor group.

4. Discussion

The decrease in activation of cell-mediated immune responses with aging may be related to the age-associated increase in tumor incidence<sup>15–17</sup>. Besides CD28 and CD95<sup>18</sup>, Treg cells play a crucial role. CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells show high expression of cell-surface molecules, including CD25, cytotoxic T-lymphocyte-associated protein-4 (CTLA4) and glucocorticoid-induced tumor necrosis factor receptor (GITR), and are characterized by expression



**Fig. 3.** Flow cytometric analysis of CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells for the six groups. Representative FACS analyses of CD4<sup>+</sup>-cell-gated spleen preparations from the six groups are presented. The top row represents the healthy group, and the bottom row the tumor group.




Fig. 4. Gene amplification for *Foxp3* was carried out successfully by RT-PCR.

of a master regulator gene, forkhead transcription factor 3 (*Foxp3*)<sup>19,20</sup>. The constitutive expression of these factors is involved in immunosuppression via several molecular mechanisms. CTLA4 interacts with CD80 and CD86 co-stimulatory molecules expressed by antigen-presenting cells (APCs) and transduces a suppressive signal<sup>21,22</sup>. Alternatively, it can directly mediate suppression by triggering induction of the enzyme indoleamine 2,3-dioxygenase in dendritic cells (DCs), which catalyzes the conversion of tryptophan to kynurenine and other metabolites and has potent immunosuppressive effects on DCs. GITR is also present on the surface of CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cells and its ligand is expressed on APCs in mice and on plasmacytoid DCs in human; much evidence indicates that it plays a role in suppressing effector T cells<sup>23,24</sup>. The

transcription factor *Foxp3* is a key player in conferring a suppressive function to CD4<sup>+</sup> T cells<sup>25,26</sup>. Retroviral transduction of *Foxp3* upregulates Treg-associated molecules, including CD25, CTLA4 and GITR, and suppresses gene transcription for cytokines such as IL-10 and TGF- $\beta$ <sup>20</sup>. Besides mechanisms that depend on cell–cell contact, CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cells are involved in the maintenance of downregulation of various immune responses by these cytokines in an indirect manner<sup>27</sup>.

In the present study, we set up a Lewis lung tumor model in mice of different ages and analyzed changes in the number of CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cells and in *Foxp3* mRNA levels in splenocytes. Our results revealed significant increases in both parameters for tumor groups compared with the corresponding healthy groups, which is consistent with previous studies<sup>28,29</sup>. In the tumor groups, as a result of immunosuppression, the increase in CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cells further supported tumor immune evasion, whereas other studies demonstrated no correlation between a higher number of CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cells and suppression activity<sup>30,31</sup>. It is not known precisely why CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cells are more numerous in mice with tumors compared to their healthy counterparts. Research indicates that natural CD4<sup>+</sup>CD25<sup>+</sup> T cells are converted to Treg cells by IL-10 and TGF- $\beta$ , which are tumor-derived soluble factors, under tumor conditions<sup>32,33</sup>.

Results for the healthy groups revealed that the number of the CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cells and *Foxp3* mRNA levels increased with aging. Despite a wealth of studies, there are no obvious explanations of why the CD4<sup>+</sup>CD25<sup>+</sup>*Foxp3*<sup>+</sup> Treg cell population




Fig. 5. *Foxp3* mRNA expression in spleen from the six groups. Data are mean  $\pm$  SD values after normalization to *HPRT* levels.



increases with aging in healthy individuals and where it comes from. Levels of both IL-10<sup>34</sup> and TGF- $\beta$ <sup>35</sup> increase with aging, and CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells in elderly mice might be generated from CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>-</sup> T cells, induced by changes in TGF- $\beta$  and IL-10 in a tumor environment. Another important explanation is the existence of two subsets of CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells with very distinct fates, as confirmed by Fisson et al.<sup>36</sup>. One subset exhibits a rapid proliferation rate characterized by a CD44<sup>high</sup> phenotype, whereas the other subset does not divide but is long-lived, and is characterized by a CD44<sup>int</sup> phenotype. These two subsets might be relevant to the increase in number, but the mechanism has not been completely clarified.

Our experimental results reveal a close relationship between changes in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells and aging and lung tumor genesis and development. CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells play an important role in inhibiting immune responses, and increases in this cell population with aging reduce the activity of the immune system. Thus, the increase in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells with aging may be a major factor in tumor incidence and the damage caused by this increase may translate into a susceptibility to tumorigenesis in the elderly. Evaluation of changes in CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> Treg cells with aging to establish a normal range for these cells at different ages could help in providing a threshold target for immune crises in the elderly and in predicting tumor occurrence. Furthermore, if questions related to the mechanism involved, such as immunological tolerance and accumulation, can be answered, it should be possible to take effective measures to interfere with this mechanism, which would open new prospects for Treg-targeted immunomodulatory therapy.

## Acknowledgments

This study was supported by the Summit on Six Aspects of Talented People and the Scientific and Technological Fund to Support the Project of Suzhou City (ZS0901).

## References

1. Anisimov VN. Carcinogenesis and aging 20 years after: escaping horizon. *Mech Aging Dev.* 2009;130:105–121.
2. Franceschi S, La Vecchia C. Cancer epidemiology in the elderly. *Crit Rev Oncol Hematol.* 2001;39:219–226.
3. Bürkle A, Caselli G, Franceschi C, et al. Pathophysiology of ageing, longevity and age related diseases. *Immun Ageing.* 2007;4:1–8.
4. Pawelec G, Solana R. Are cancer and ageing different sides of the same coin? Conference on cancer and ageing. *EMBO Rep.* 2008;9:234–238.
5. Campisi J, Yaswen P. Aging and cancer cell biology. *Aging Cell.* 2009;8:221–225.
6. Irmiger-Finger I. Science of cancer and aging. *J Clin Oncol.* 2007;25:1844–1851.
7. Bhardwaj N. Harnessing the immune system to treat cancer. *J Clin Invest.* 2007;117:1130–1136.
8. Piccirillo CA, Shevach EM. Cutting edge: control of CD8<sup>+</sup> T cell activation by CD4<sup>+</sup>CD25<sup>+</sup> immunoregulatory cells. *J Immunol.* 2001;167:1137–1140.
9. Suri-Payer E, Amar AZ, Thornton AM, et al. CD4<sup>+</sup>CD25<sup>+</sup> T cells inhibit both the induction and effector function of autoreactive T cells and represent a unique lineage of immunoregulatory cells. *J Immunol.* 1998;160:1212–1218.
10. Shevach EM, McHugh RS, Piccirillo CA, et al. Control of T-cell activation by CD4<sup>+</sup>CD25<sup>+</sup> suppressor T cells. *Immunol Rev.* 2001;182:58–67.
11. Zou WP. Regulatory T cells, tumor immunity and immunotherapy. *Nat Rev Immunol.* 2006;6:295–307.
12. Gregg R, Smith CM, Clark FJ, et al. The number of human peripheral blood CD4<sup>+</sup>CD25<sup>high</sup> regulatory T cells increases with age. *Clin Exp Immunol.* 2005;140:540–546.
13. Parkin DM, Bray FB, Pisani P. Global cancer statistics. *CA Cancer J Clin.* 2002;2005(55):74–108.
14. Batura-Gabryel H, Foremska-Iciek J. Lung cancer in the elderly – increasing epidemiological problem of 21st century. *Rocz Akad Med Bialymst.* 2005;50:152–155.
15. Provinciali M, Smorlesi A. Immunoprevention and immunotherapy of cancer in aging. *Cancer Immunol Immunother.* 2005;54:93–106.
16. Linton P, Thoman ML. T cell senescence. *Front Biosci.* 2001;6:248–261.
17. Lustgarten J, Dominguez AL, Thoman M. Aged mice develop protective anti-tumor immune responses with appropriate costimulation. *J Immunol.* 2004;173:4510–4515.
18. Wang L, Pan XD, Xie Y, et al. Altered CD28 and CD95 mRNA expression in peripheral blood mononuclear cells from elderly patients with primary non-small cell lung cancer. *Chin Med J.* 2010;123:51–56.
19. Yagi H, Nomura T, Nakamura K, et al. Crucial role of FOXP3 in the development and function of human CD25<sup>+</sup>CD4<sup>+</sup> regulatory T cells. *Int Immunol.* 2004;16:1643–1656.
20. Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science.* 2003;299:1057–1061.
21. Sakaguchi S. Naturally arising CD4<sup>+</sup> regulatory T cells for immunologic self-tolerance and negative control of immune responses. *Annu Rev Immunol.* 2004;22:531–562.
22. Huang CT, Workman CJ, Flies D, et al. Role of LAG-3 in regulatory T cells. *Immunity.* 2004;21:503–513.
23. Shimizu J, Yamazaki S, Takahashi T, et al. Stimulation of CD25<sup>+</sup>CD4<sup>+</sup> regulatory T cells through GITR breaks immunological self-tolerance. *Nat Immunol.* 2002;3:135–142.
24. Shevach EM, Stephens GL. The GITR–GITRL interaction: co-stimulation or contrasuppression of regulatory activity? *Nat Rev Immunol.* 2006;6:613–618.
25. Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4<sup>+</sup>CD25<sup>+</sup> regulatory T cells. *Nat Immunol.* 2003;4:330–336.
26. Khattry R, Cox T, Yasayko SA, et al. An essential role for scurf in CD4<sup>+</sup>CD25<sup>+</sup> T regulatory cells. *Nat Immunol.* 2003;4:337–342.
27. Muller AJ, Scherle PA. Targeting the mechanisms of tumoral immune tolerance with small-molecule inhibitors. *Nat Rev Cancer.* 2006;6:613–625.
28. Sharma S, Dominguez AL, Lustgarten J. High accumulation of T regulatory cells prevents the activation of immune responses in aged animals. *J Immunol.* 2006;177:8348–8355.
29. Zhao L, Sun L, Wang H, et al. Changes of CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> regulatory T cells in aged BALB/c mice. *J Leukocyte Biol.* 2007;81:1386–1394.
30. Beyer M, Kochanek M, Darabi K, et al. Reduced frequencies and suppressive function of CD4<sup>+</sup>CD25<sup>hi</sup> regulatory T cells in patients with chronic lymphocytic leukemia after therapy with fludarabine. *Blood.* 2005;106:2018–2025.
31. Putheti P, Pettersson A, Soderstrom M, et al. Circulating CD4<sup>+</sup>CD25<sup>+</sup> T regulatory cells are not altered in multiple sclerosis and unaffected by disease-modulating drugs. *J Clin Immunol.* 2004;24:155–161.
32. Moo-Young TA, Larson JW, Belt BA, et al. Tumor-derived TGF-beta mediates conversion of CD4<sup>+</sup>Foxp3<sup>+</sup> regulatory T cells in a murine model of pancreas cancer. *J Immunother.* 2009;32:12–21.
33. Chen W, Jin W, Hardegen N, et al. Conversion of peripheral CD4<sup>+</sup>CD25<sup>-</sup> naive T cells to CD4<sup>+</sup>CD25<sup>+</sup> regulatory T cells by TGF-beta induction of transcription factor Foxp3. *J Exp Med.* 2003;198:1875–1886.
34. Pioli C, Pucci S, Barile S, et al. Role of mRNA stability in the different patterns of cytokine production by CD4<sup>+</sup> cells from young and old mice. *Immunology.* 1998;94:380–387.
35. Wan YY, Flavell RA. The roles for cytokines in the generation and maintenance of regulatory T cells. *Immunol Rev.* 2006;212:114–130.
36. Fisson S, Darrasse-Jèze G, Litvinova E, et al. Continuous activation of autoreactive CD4<sup>+</sup>CD25<sup>+</sup> regulatory T cells in the steady state. *J Exp Med.* 2003;198:737–746.